

Aspects of domain-wall fermions on the lattice

Ting-Wai Chiu

Department of Physics, National Taiwan University

Taipei, Taiwan 106, Taiwan.

E-mail : *twchiu@phys.ntu.edu.tw*

Abstract

I introduce two new boundary layers to the optimal domain-wall fermion action, and formulate vector gauge theories (e.g., QCD) with the new action. The generating functional for n -point Green's function of quark fields is derived. The bilinear and higher-order quark observables are shown to be a_5 -invariant, to possess optimal chiral symmetry for any finite N_s , as well as to preserve continuum discrete symmetries at finite lattice spacing. With 2D abelian background gauge fields, I compute the index, the axial anomaly, and the fermion determinant (ratio) of the effective lattice Dirac operator, for the optimal DWF and the conventional DWF respectively. All quantities of the optimal DWF are in good agreement with the exact solution, but the fermion determinant (ratio) of the conventional DWF disagrees with the exact solution, for topologically nontrivial sectors.

PACS numbers: 11.15.Ha, 11.30.Rd, 12.38.Gc

Keywords : Lattice QCD, Domain-wall fermions, Overlap Dirac operator, Zolotarev optimal rational approximation.

1 Introduction

The basic idea of domain-wall fermions (DWF) [1, 2] is to use an infinite set of coupled Dirac fermion fields [$\psi_s(x), s \in (-\infty, \infty)$] with masses behaving like a step function $m(s) = m\theta(s)$ such that Weyl fermion states can arise as zeromodes bound to the mass defect at $s = 0$. However, if one uses a compact set of masses, then the boundary conditions of the mass (step) function must lead to the occurrence of both left-handed and right-handed chiral fermion fields, i.e., a vector-like theory. For lattice QCD with DWF [3], in practice, one can only use a finite number (N_s) of lattice Dirac fermion fields to set up the domain wall, thus the chiral symmetry of the quark fields (in the massless limit) is broken. Therefore, a relevant question is how to construct a domain-wall fermion action such that the effective 4D lattice Dirac operator has the optimal chiral symmetry for any finite N_s . Nevertheless, the chiral symmetry is not the only concern of lattice QCD with domain-wall fermions. Another crucial question is whether the effective 4D lattice Dirac operator depends on the lattice spacing a_5 in the fifth dimension. If the dependence on a_5 is strong, it is unclear what is the proper value of a_5 for measuring physical observables, assuming N_s and the lattice spacing a are fixed.

Recently, a lattice domain-wall fermion action has been constructed [5] such that the effective 4D lattice Dirac operator preserves the chiral symmetry optimally for any finite N_s , and is invariant for any a_5 . Further, it has been asserted that the effective 4D lattice Dirac operator for the internal fermion loops is exponentially-local for sufficiently smooth gauge backgrounds [6].

Explicitly, the action of optimal lattice domain-wall fermions reads¹

$$\begin{aligned} \mathcal{A}_f &= \sum_{s,s'=1}^{N_s} \sum_{x,x'} \bar{\psi}(x,s) \{ (\omega_s a_5 D_w(x,x') + \delta_{x,x'}) \delta_{s,s'} \\ &\quad - (\delta_{x,x'} - \omega_s a_5 D_w(x,x')) (P_- \delta_{s',s+1} + P_+ \delta_{s',s-1}) \} \psi(x',s') \\ &\equiv \sum_{s,s'=1}^{N_s} \sum_{x,x'} \bar{\psi}(x,s) \mathcal{D}(x,s;x's') \psi(x',s'), \quad P_{\pm} = \frac{1 \pm \gamma_5}{2} \end{aligned} \quad (1)$$

with boundary conditions

$$P_+ \psi(x, 0) = -r m_q P_+ \psi(x, N_s), \quad r = \frac{1}{2m_0} \quad (2)$$

$$P_- \psi(x, N_s + 1) = -r m_q P_- \psi(x, 1), \quad (3)$$

where m_q is the bare quark mass, D_w is the standard 4D Wilson-Dirac operator plus a negative parameter,

$$D_w = \sum_{\mu=1}^4 \gamma_{\mu} t_{\mu} + W - m_0, \quad m_0 \in (0, 2) \quad (4)$$

¹In this paper, we suppress the lattice spacing a , as well as the Dirac and color indices, which can be restored easily.

$$t_\mu(x, x') = \frac{1}{2} \left[U_\mu(x) \delta_{x', x+\mu} - U_\mu^\dagger(x') \delta_{x', x-\mu} \right], \quad (5)$$

$$W(x, x') = \sum_{\mu=1}^4 \frac{1}{2} \left[2\delta_{x, x'} - U_\mu(x) \delta_{x', x+\mu} - U_\mu^\dagger(x') \delta_{x', x-\mu} \right], \quad (6)$$

and the weights

$$\omega_s a_5 = \frac{1}{\lambda_{min}} \sqrt{1 - \kappa'^2 \text{sn}^2(v_s; \kappa')}, \quad s = 1, \dots, N_s. \quad (7)$$

Here $\text{sn}(v_s; \kappa')$ is the Jacobian elliptic function with argument v_s (12) and modulus $\kappa' = \sqrt{1 - \lambda_{min}^2 / \lambda_{max}^2}$, where λ_{max} and λ_{min} are the maximum and the minimum of the eigenvalues of $|H_w|$,

$$H_w = \gamma_5 D_w. \quad (8)$$

The weights in (7) are obtained from the roots ($u_s = (\omega_s a_5)^{-2}$, $s = 1, \dots, N_s$) of the equation

$$\delta_Z(u) = \begin{cases} 1 - \sqrt{u} R_Z^{(n,n)}(u) = 0, & N_s = 2n + 1 \\ 1 - \sqrt{u} R_Z^{(n-1,n)}(u) = 0, & N_s = 2n \end{cases}$$

such that the operator S_{opt} (22) in the effective 4D lattice Dirac operator is equal to the optimal rational approximation of $\text{sgn}(H_w) = H_w (H_w^2)^{-1/2}$,

$$S_{opt} = S_{opt}(H_w) = \begin{cases} H_w R_Z^{(n,n)}(H_w^2), & N_s = 2n + 1, \\ H_w R_Z^{(n-1,n)}(H_w^2), & N_s = 2n. \end{cases} \quad (9)$$

where $R_Z(H_w^2)$ is the Zolotarev optimal rational polynomial [7, 8] for the inverse square root of H_w^2 ,

$$R_Z^{(n,n)}(H_w^2) = \frac{d_0}{\lambda_{min}} \prod_{l=1}^n \frac{1 + h_w^2 / c_{2l}}{1 + h_w^2 / c_{2l-1}}, \quad h_w^2 = H_w^2 / \lambda_{min}^2 \quad (10)$$

and

$$R_Z^{(n-1,n)}(H_w^2) = \frac{d'_0}{\lambda_{min}} \frac{\prod_{l=1}^{n-1} (1 + h_w^2 / c'_{2l})}{\prod_{l=1}^n (1 + h_w^2 / c'_{2l-1})}, \quad (11)$$

and the coefficients d_0 , d'_0 , c_l and c'_l are expressed in terms of elliptic functions [8] with arguments depending only on N_s and $\lambda_{max}^2 / \lambda_{min}^2$.

The argument v_s in (7) is

$$v_s = (-1)^{s-1} M \text{sn}^{-1} \left(\sqrt{\frac{1 + 3\lambda}{(1 + \lambda)^3}}; \sqrt{1 - \lambda^2} \right) + \left[\frac{s}{2} \right] \frac{2K'}{N_s} \quad (12)$$

where

$$\lambda = \prod_{l=1}^{N_s} \frac{\Theta^2\left(\frac{2lK'}{N_s}; \kappa'\right)}{\Theta^2\left(\frac{(2l-1)K'}{N_s}; \kappa'\right)}, \quad (13)$$

$$M = \prod_{l=1}^{\lfloor \frac{N_s}{2} \rfloor} \frac{\text{sn}^2\left(\frac{(2l-1)K'}{N_s}; \kappa'\right)}{\text{sn}^2\left(\frac{2lK'}{N_s}; \kappa'\right)}, \quad (14)$$

K' is the complete elliptic integral of the first kind with modulus κ' , and Θ is the elliptic theta function. From (7), it is clear that $\lambda_{max}^{-1} \leq a_5 \omega_s \leq \lambda_{min}^{-1}$, since $\text{sn}^2(\cdot) \leq 1$.

Obviously, the optimal domain-wall fermion action (1) is invariant for any a_5 , since its dependence on a_5 only through the product $\omega_s a_5$ (7) which only depends on N_s , λ_{min} and $\lambda_{max}^2/\lambda_{min}^2$. Therefore, a_5 is a *redundant parameter in the optimal domain-wall fermion action* (1).

Now the effective 4D lattice Dirac operator of the optimal domain-wall fermion action (1) can be extracted from its fermion determinant

$$\det \mathcal{D}(m_q) = \int [d\bar{\psi}][d\psi] e^{-\mathcal{A}_f[\psi, \bar{\psi}]}, \quad (15)$$

where $\mathcal{D}(m_q)$ is the 5-dimensional optimal DWF operator defined in (1). The Grassman integral on the r.h.s. of (15) can be evaluated by transforming $\{\psi_s, \bar{\psi}_s\}$ to $\{\eta_s, \bar{\eta}_s\}$ according to

$$\psi_1 = P_- \eta_1 + P_+ \eta_2, \quad \bar{\psi}_1 = \bar{\eta}_1 (\omega_1 a_5 H_w - 1)^{-1} \gamma_5, \quad (16)$$

$$\psi_s = P_- \eta_s + P_+ \eta_{s+1}, \quad \bar{\psi}_s = \bar{\eta}_s (\omega_s a_5 H_w - 1)^{-1} \gamma_5, \quad s = 2, \dots, N_s - 1 \quad (17)$$

$$\psi_{N_s} = P_- \eta_{N_s} + P_+ \eta_1, \quad \bar{\psi}_{N_s} = \bar{\eta}_{N_s} (\omega_{N_s} a_5 H_w - 1)^{-1} \gamma_5, \quad (18)$$

and then integrating $\{\eta_s, \bar{\eta}_s\}$ successively from $s = N_s$ to $s = 1$. The result (will be derived in Section 2) is

$$\det \mathcal{D}(m_q) = K \det[r(D_c + m_q)] \quad (19)$$

where

$$K = \prod_{s=1}^{N_s} \det[\gamma_5 (\omega_s a_5 H_w - 1)] \det\left(-P_+ + \prod_{s=1}^{N_s} T_s^{-1} P_-\right), \quad (20)$$

$$r D_c = \frac{1 + \gamma_5 S_{opt}}{1 - \gamma_5 S_{opt}}, \quad (21)$$

$$S_{opt} = \frac{1 - \prod_{s=1}^{N_s} T_s}{1 + \prod_{s=1}^{N_s} T_s}, \quad (22)$$

$$T_s = \frac{1 - \omega_s a_5 H_w}{1 + \omega_s a_5 H_w}, \quad s = 1, \dots, N_s. \quad (23)$$

Even though D_c is chirally symmetric in the limit $N_s \rightarrow \infty$, one still cannot identify $(D_c + m_q)$ as the effective 4D lattice Dirac operator for the internal quark fields, since D_c is nonlocal and $\det(D_c + m_q)$ does not reproduce $\det[\gamma_\mu(\partial_\mu + iA_\mu) + m_q]$ in the continuum limit ($a \rightarrow 0$). (It is understood that the fermion determinant is normalized by its free fermion limit.) Also, one cannot neglect the factor K which can be written as the determinant of a 4D lattice fermion operator.

The resolution is to introduce a set of pseudofermions (Pauli-Villars) fields $\{\phi, \bar{\phi}\}$ similar to the domain-wall fermions but obeying Bose statistics. Explicitly, the action of the pseudofermion fields reads

$$\begin{aligned} \mathcal{A}_{pf} = & \sum_{s,s'=1}^{N_s} \sum_{x,x'} \bar{\phi}(x, s) \{ (\omega_s a_5 D_w(x, x') + \delta_{x,x'}) \delta_{s,s'} \\ & - (\delta_{x,x'} - \omega_s a_5 D_w(x, x')) (P_- \delta_{s',s+1} + P_+ \delta_{s',s-1}) \} \phi(x', s') \end{aligned} \quad (24)$$

with boundary conditions

$$P_+ \phi(x, 0) = -P_+ \phi(x, N_s), \quad P_- \phi(x, N_s + 1) = -P_- \phi(x, 1),$$

where $m_q = 2m_0$ has been imposed. Then, we have

$$\begin{aligned} \det[\mathcal{D}(m_q) \mathcal{D}^{-1}(2m_0)] &= \int [d\bar{\psi}][d\psi][d\bar{\phi}][d\phi] e^{-\mathcal{A}_f[\psi, \bar{\psi}] - \mathcal{A}_{pf}[\phi, \bar{\phi}]} \\ &= \det[r(D_c + m_q)(1 + rD_c)^{-1}]. \end{aligned}$$

Thus the effective 4D lattice Dirac operator for the internal quark fields (dressed with pseudofermions) can be identified as

$$D(m_q) = (D_c + m_q)(1 + rD_c)^{-1} = m_q + (m_0 - m_q/2) (1 + \gamma_5 H_w R_Z(H_w)) \quad (25)$$

where $R_Z(H_w^2)$ is the Zolotarev rational polynomial defined in (9)-(11). The exponential locality of $D(m_q)$ (for any m_q and N_s) has been asserted for sufficiently smooth gauge backgrounds [6].

In the limit $N_s \rightarrow \infty$, S_{opt} becomes the sign function of H_w

$$\lim_{N_s \rightarrow \infty} S_{opt} = \frac{H_w}{\sqrt{H_w^2}},$$

and D_c (21) is exactly chirally symmetric

$$\lim_{N_s \rightarrow \infty} (D_c \gamma_5 + \gamma_5 D_c) = 0, \quad (26)$$

which is equivalent to

$$\lim_{\substack{N_s \rightarrow \infty \\ m_q \rightarrow 0}} [(D_c + m_q)^{-1} \gamma_5 + \gamma_5 (D_c + m_q)^{-1}] = 0, \quad (27)$$

the chiral symmetry of the valence quark propagator $(D_c + m_q)^{-1}$ in the massless limit. Therefore one can use this chirally symmetric (but nonlocal) D_c to construct an exponentially-local lattice Dirac operator [9]

$$D = D_c(1 + rD_c)^{-1} \quad (28)$$

which satisfies the Ginsparg-Wilson relation [10]

$$D\gamma_5 + \gamma_5 D = 2rD\gamma_5 D, \quad (29)$$

and its massive generalization [11]

$$D(m) = (D_c + m)(1 + rD_c)^{-1}, \quad (30)$$

which is exactly in the same form of the effective 4D lattice Dirac operator (25) for any N_s .

Obviously, the optimal DWF action (1) in the limit $N_s \rightarrow \infty$ realizes the exact chiral symmetry (27) as well as the Ginsparg-Wilson relation (29). Further, for any finite N_s , the action (1) preserves the chiral symmetry optimally since the deviation of S_{opt} from the sign function is the minimum, and has a theoretical upper bound [12],

$$\Delta_Z = \max_{\forall Y \neq 0} \left| \frac{Y^\dagger S_{opt}^2 Y}{Y^\dagger Y} - 1 \right| \leq \frac{2(1 - \lambda)}{1 + \lambda} \equiv B_Z(N_s, b) \quad (31)$$

where λ is defined in (13), a function of N_s and $b = \lambda_{max}^2 / \lambda_{min}^2$. Thus, for any given gauge configuration, one can use the theoretical error bound to determine what values of N_s and b (i.e., how many low-lying eigenmodes of H_w^2 should be projected out) are required to attain one's desired accuracy in preserving the exact chiral symmetry. In practice, with $N_s = 32$, one should have no difficulties to achieve $\Delta_Z < 10^{-12}$ for any gauge configurations on a finite lattice (say, $16^3 \times 32$ at $\beta = 6.0$). The theoretical error bound can be estimated as

$$B_Z(N_s, b) = A(b)e^{-c(b)N_s}$$

where

$$\begin{aligned} c(b) &\simeq 4.27(45) \ln(b)^{-0.746(5)}, \\ A(b) &\simeq 8.11(1) b^{-0.0091(1)} \ln(b)^{0.0042(3)}. \end{aligned}$$

Note that D (28) ($N_s \rightarrow \infty$) times a factor $r = 1/2m_0$ is exactly equal to the overlap Dirac operator [13, 14],

$$D_o = \frac{1}{2} \left(1 + \gamma_5 \frac{H_w}{\sqrt{H_w^2}} \right). \quad (32)$$

This implies that D is topologically-proper (i.e., with the correct index and axial anomaly), similar to the case of overlap Dirac operator. For any finite N_s , r times $D(m_q = 0)$ (25) is exactly equal to the overlap Dirac operator D_o with $(H_w^2)^{-1/2}$ approximated by Zolotarev optimal rational polynomial, (10) ($N_s = \text{odd}$), or (11) ($N_s = \text{even}$).

Although the effective 4D lattice Dirac operator of the optimal DWF action is a_5 -invariant and preserves optimal chiral symmetry for any finite N_s , the correlation function of quark fields $q(x)$ and $\bar{q}(y)$ is not equal to the quark propagator $(D_c + m_q)_{x,y}^{-1}$, nor the inverse of the effective 4D lattice Dirac operator, $D(m_q)_{x,y}^{-1}$. This is unsatisfactory from the theoretical viewpoint. In this paper, I introduce two new boundary layers to the optimal DWF action, and derive the generating functional for n -point Green's function of quark fields q and $\bar{q}$. Then the bilinear (and higher-order) quark operators manifest their continuum discrete symmetries, besides the a_5 -invariance and optimal chiral symmetry for any finite N_s . With 2D abelian background gauge fields, I compute the index, the axial anomaly, and the fermion determinant (ratio) of the effective lattice Dirac operator, for the optimal DWF and the conventional DWF respectively. All quantities of the optimal DWF are in good agreement with the exact solution, but the fermion determinant (ratio) of the conventional DWF disagrees with the exact solution, for topologically nontrivial sectors.

The outline of this paper is as follows. In Section 2, we derive the generating functional for n -point Green's function of quarks fields in the optimal DWF as well as the conventional DWF, and discuss the issues pertinent to finite N_s and a_5 . The discrete symmetry (V-A structure) of the left-handed quark current in effective weak Hamiltonian is shown explicitly. In Section 3, we compute the index, the axial anomaly, and the fermion determinant of the effective lattice Dirac operators in 2D abelian background gauge fields. In Section 4, we summarize and conclude with a discussion on the difference between the realization of exact chiral symmetry via the domain-wall fermions and the Ginsparg-Wilson lattice Dirac operator. In the Appendix, we derive a formula for computing the propagator of the effective 4D lattice Dirac operator (25) of the optimal DWF (1).

2 Generating functional for n -point Green's function

In quantum field theory, one of the fundamental quantities is the generating functional $Z[J]$ ($W[J] = \ln Z[J]$) for (connected) n -point Green's function. Once it is derived, one can obtain any (connected) n -point Green's function by successive differentiations with respect to the source J . Further, the generating functional for one-particle irreducible vertices, $\Gamma[\phi]$, can also be obtained

through the Legendre transform $\Gamma[\phi] = W[J] - \int dx J(x)\phi(x)$. In this Section, we derive the generating functional $Z[J] = e^{W[J]}$ for n -point Green's function of quark fields q and $\bar{q}$ constructed from the boundary modes of domain-wall fermions.

The generating functional for n -point Green's function of quark q and $\bar{q}$ is defined as

$$e^{W[J, \bar{J}]} = Z[J, \bar{J}] = \frac{\int [dU][d\bar{\psi}][d\psi][d\bar{\phi}][d\phi] e^{-\mathcal{A}_g - \mathcal{A}_f - \mathcal{A}_{pf} + \sum_x \{ \bar{J}(x)q(x) + \bar{q}(x)J(x) \}}}{\int [dU][d\bar{\psi}][d\psi][d\bar{\phi}][d\phi] e^{-\mathcal{A}_g - \mathcal{A}_f - \mathcal{A}_{pf}}} \quad (33)$$

where $\mathcal{A}_g$ is the gauge field action, $\mathcal{A}_f$ is the domain-wall fermion action, $\mathcal{A}_{pf}$ is the corresponding pseudofermion action, and $\bar{J}$ and J are the Grassman sources of q and $\bar{q}$ respectively.

We first derive $Z[J, \bar{J}] = e^{W[J, \bar{J}]}$ for the optimal DWF action, then show that it is invariant for any a_5 , and preserves the chiral symmetry optimally for any finite N_s . Then we derive $Z[J, \bar{J}]$ for the usual DWF action, and discuss its problems due to finite N_s and a_5 .

2.1 Optimal domain-wall fermions

First, I generalize the optimal domain fermion action (1) by appending two boundary layers at $s = 0$ and $s = N_s + 1$,

$$\begin{aligned} \mathcal{A}'_f &= \sum_{s, s'=0}^{N_s+1} \sum_{x, x'} \bar{\psi}(x, s) \{ (\omega_s a_5 D_w(x, x') + \delta_{x, x'}) \delta_{s, s'} \\ &\quad - (\delta_{x, x'} - \omega_s a_5 D_w(x, x')) (P_- \delta_{s', s+1} + P_+ \delta_{s', s-1}) \} \psi(x', s') \\ &\equiv \sum_{s, s'=0}^{N_s+1} \sum_{x, x'} \bar{\psi}(x, s) \mathcal{D}'(x, s; x', s') \psi(x', s') \end{aligned} \quad (34)$$

with boundary conditions

$$P_+ \psi(x, -1) = -rm_q P_+ \psi(x, N_s + 1), \quad r = \frac{1}{2m_0} \quad (35)$$

$$P_- \psi(x, N_s + 2) = -rm_q P_- \psi(x, 0), \quad (36)$$

where the weights at the boundaries are zero, i.e. $\omega_0 = \omega_{N_s+1} = 0$, and other weights are given by (7) as before.

The quark fields coupling to physical hadrons are constructed from the left and right boundary modes

$$q(x) = P_- \psi(x, 0) + P_+ \psi(x, N_s + 1), \quad (37)$$

$$\bar{q}(x) = \bar{\psi}(x, 0) P_+ + \bar{\psi}(x, N_s + 1) P_- . \quad (38)$$

It will be shown that the fermion determinant of the new action (34) is exactly equal to that (19) of the original action (1). Thus the effective 4D

lattice Dirac operator is exactly equal to (25), after the pseudofermion action (with $m_q = 2m_0$) is incorporated. The purpose of adding two boundary layers at $s = 0$ and $s = N_s + 1$ is for the construction of the quark fields (37)-(38) such that the correlation function of $q(x)$ and $\bar{q}(y)$ is equal to the valence quark propagator, i.e., $r\langle q(x)\bar{q}(y) \rangle = (D_c + m_q)_{x,y}^{-1}$ in any background gauge field, and the generating functional $Z[J, \bar{J}]$ can be derived.

Now we derive the generating functional $Z[J, \bar{J}]$ for the optimal DWF (34), according to the formula (33).

Using $\gamma_5 P_{\pm} = \pm P_{\pm}$, $P_+ + P_- = 1$, and $H_w = \gamma_5 D_w$, we can rewrite (34) as

$$\begin{aligned} \mathcal{A}'_f = & (rm_q + 1)\bar{\psi}_0\gamma_5 P_+ \psi_{N_s+1} \\ & - \bar{\psi}_0\gamma_5(P_- \psi_0 + P_+ \psi_{N_s+1}) + \bar{\psi}_0\gamma_5(P_- \psi_1 + P_+ \psi_0) \\ & + \sum_{s=1}^{N_s} \left\{ \bar{\psi}_s\gamma_5(\omega_s a_5 H_w - 1)(P_- \psi_s + P_+ \psi_{s-1}) + \bar{\psi}_s\gamma_5(\omega_s a_5 H_w + 1)(P_- \psi_{s+1} + P_+ \psi_s) \right\} \\ & - \bar{\psi}_{N_s+1}\gamma_5(P_- \psi_{N_s+1} + P_+ \psi_{N_s}) + \bar{\psi}_{N_s+1}\gamma_5(P_- \psi_0 + P_+ \psi_{N_s+1}) \\ & - (rm_q + 1)\bar{\psi}_{N_s+1}\gamma_5 P_- \psi_0 \end{aligned} \quad (39)$$

where all indices are suppressed except the flavor index in the fifth dimension.

Next we define

$$\eta_0 = P_- \psi_0 + P_+ \psi_{N_s+1}, \quad \bar{\eta}_0 = -\bar{\psi}_0\gamma_5, \quad (40)$$

$$\eta_s = P_- \psi_s + P_+ \psi_{s-1}, \quad \bar{\eta}_s = \bar{\psi}_s\gamma_5(\omega_s a_5 H_w - 1), \quad s = 1, \dots, N_s \quad (41)$$

$$\eta_{N_s+1} = P_- \psi_{N_s+1} + P_+ \psi_{N_s}, \quad \bar{\eta}_{N_s+1} = -\bar{\psi}_{N_s+1}\gamma_5, \quad (42)$$

and the inverse transform

$$\psi_0 = P_- \eta_0 + P_+ \eta_1, \quad \bar{\psi}_0 = -\bar{\eta}_0\gamma_5, \quad (43)$$

$$\psi_s = P_- \eta_s + P_+ \eta_{s+1}, \quad \bar{\psi}_s = \bar{\eta}_s(\omega_s a_5 H_w - 1)^{-1}\gamma_5, \quad s = 1, \dots, N_s \quad (44)$$

$$\psi_{N_s+1} = P_- \eta_{N_s+1} + P_+ \eta_0, \quad \bar{\psi}_{N_s+1} = -\bar{\eta}_{N_s+1}\gamma_5. \quad (45)$$

Then the action (39) can be expressed in terms of $\{\eta, \bar{\eta}\}$ fields

$$\begin{aligned} \mathcal{A}'_f = & \bar{\eta}_0(P_- - rm_q P_+)\eta_0 - \bar{\eta}_0\eta_1 + \sum_{s=1}^{N_s} \left\{ \bar{\eta}_s\eta_s - \bar{\eta}_s T_s^{-1}\eta_{s+1} \right\} \\ & + \bar{\eta}_{N_s+1}\eta_{N_s+1} - \bar{\eta}_{N_s+1}(P_+ - rm_q P_-)\eta_0 \end{aligned} \quad (46)$$

where

$$T_s = \frac{1 - \omega_s a_5 H_w}{1 + \omega_s a_5 H_w}, \quad s = 1, \dots, N_s.$$

Thus the fermionic integral in the numerator of (33) can be written as

$$\int [d\bar{\psi}][d\psi] e^{-\mathcal{A}'_f[\psi, \bar{\psi}] + \bar{J}q + \bar{q}J} = \mathcal{J} \int [d\bar{\eta}][d\eta] e^{-\mathcal{A}'_f[\eta, \bar{\eta}] + \bar{J}\eta_0 - \bar{\eta}_0 P_+ J + \bar{\eta}_{N_s+1} P_- J} \quad (47)$$

where

$$\begin{aligned}\bar{J}q + \bar{q}J &= \bar{J}(P_- \psi_0 + P_+ \psi_{N_s+1}) + (\bar{\psi}_0 P_+ + \bar{\psi}_{N_s+1} P_-)J \\ &= \bar{J}\eta_0 - \bar{\eta}_0 P_+ J + \bar{\eta}_{N_s+1} P_- J\end{aligned}\quad (48)$$

and $\mathcal{J}$ is the Jacobian of the transformation,

$$\mathcal{J} = \prod_{s=0}^{N_s+1} \det [\gamma_5(\omega_s a_5 H_w - 1)] = \prod_{s=1}^{N_s} \det [\gamma_5(\omega_s a_5 H_w - 1)] \quad (49)$$

where $\omega_0 = \omega_{N_s+1} = 0$ has been used.

Now using the Grassman integral formula

$$\int d\bar{\chi} d\chi e^{-\bar{\chi} M \chi + \bar{v} \chi + \bar{\chi} v} = e^{\bar{v} M^{-1} v} \det M \quad (50)$$

one can easily evaluate the Grassman integrals in (47), by integrating $(\eta_s, \bar{\eta}_s)$ successively from $s = N_s + 1$ to $s = 0$.

Explicitly, after integrating $(\eta_{N_s+1}, \bar{\eta}_{N_s+1})$, (47) becomes

$$\begin{aligned}\mathcal{J} \int \prod_{s=0}^{N_s} [d\bar{\eta}_s][d\eta_s] \exp \Big\{ & -\bar{\eta}_0(P_- - rm_q P_+) \eta_0 + \bar{\eta}_0 \eta_1 + \bar{J} \eta_0 - \bar{\eta}_0 P_+ J \\ & - \sum_{s=1}^{N_s-1} (\bar{\eta}_s \eta_s - \bar{\eta}_s T_s^{-1} \eta_{s+1}) - \bar{\eta}_{N_s} \eta_{N_s} \\ & + \bar{\eta}_{N_s} T_{N_s}^{-1} [(P_+ - rm_q P_-) \eta_0 + P_- J] \Big\}\end{aligned}$$

then integrating $(\eta_{N_s}, \bar{\eta}_{N_s})$, the remaining integral becomes

$$\begin{aligned}\mathcal{J} \int \prod_{s=0}^{N_s-1} [d\bar{\eta}_s][d\eta_s] \exp \Big\{ & -\bar{\eta}_0(P_- - rm_q P_+) \eta_0 + \bar{\eta}_0 \eta_1 + \bar{J} \eta_0 - \bar{\eta}_0 P_+ J \\ & - \sum_{s=1}^{N_s-2} (\bar{\eta}_s \eta_s - \bar{\eta}_s T_s^{-1} \eta_{s+1}) - \bar{\eta}_{N_s-1} \eta_{N_s-1} \\ & + \bar{\eta}_{N_s-1} T_{N_s-1}^{-1} T_{N_s}^{-1} [(P_+ - rm_q P_-) \eta_0 + P_- J] \Big\}\end{aligned}$$

Subsequent integrations over $(\eta_{N_s-2}, \bar{\eta}_{N_s-2})$ up to $(\eta_1, \bar{\eta}_1)$ are similar to the above integration, and the result is

$$\begin{aligned}\mathcal{J} \int [d\bar{\eta}_0][d\eta_0] \exp \Big\{ & -\bar{\eta}_0(P_- - rm_q P_+) \eta_0 + \bar{J} \eta_0 - \bar{\eta}_0 P_+ J \\ & + \bar{\eta}_0 \prod_{s=1}^{N_s} T_s^{-1} [(P_+ - rm_q P_-) \eta_0 + P_- J] \Big\}\end{aligned}$$

Finally, integrating $(\eta_0, \bar{\eta}_0)$ gives

$$\begin{aligned} & \mathcal{J} \det \left[(P_- - rm_q P_+) - \prod_{s=1}^{N_s} T_s^{-1} (P_+ - rm_q P_-) \right] \times \\ & \exp \left\{ \bar{J} \left[\left(-P_+ + \prod_{s=1}^{N_s} T_s^{-1} P_- \right)^{-1} \left(P_- - \prod_{s=1}^{N_s} T_s^{-1} P_+ \right) + rm_q \right]^{-1} J \right\} \\ & = K \det[r(D_c + m_q)] \exp \left\{ \bar{J} r^{-1} (D_c + m_q)^{-1} J \right\} \end{aligned} \quad (51)$$

where the simple identity

$$\left(-P_+ + \prod_{s=1}^{N_s} T_s^{-1} P_- \right)^{-1} \left(P_- - \prod_{s=1}^{N_s} T_s^{-1} P_+ \right) = \frac{1 + \gamma_5 S_{opt}}{1 - \gamma_5 S_{opt}} = r D_c \quad (52)$$

has been used, and K , S_{opt} and $r D_c$ are defined in (20), (22) and (21) respectively. Note that any two transfer matrices commute with each other ($T_s T_{s'} = T_{s'} T_s$), thus D_c is invariant for swapping any two weights in $\{\omega_1, \dots, \omega_{N_s}\}$.

Now setting $\bar{J} = J = 0$ in (51), we obtain the fermion determinant of the optimal DWF action (34)

$$\det \mathcal{D}'(m_q) = K \det[r(D_c + m_q)] \quad (53)$$

where $\mathcal{D}'(m_q)$ is the 5-dimensional lattice DWF operator defined in (34), while $r D_c$ is the 4-dimensional lattice Dirac operator defined in (21), which becomes chirally symmetric in the limit $N_s \rightarrow \infty$, the salient feature of domain-wall fermions.

Obviously, if one repeats above steps to derive the fermion determinant of the original optimal DWF action (1), then the result is exactly equal to (53). The vital observation is that the transfer matrix T_s at $s = 0$ and $s = N_s + 1$ is equal to the identity (since $\omega_0 = \omega_{N_s+1} = 0$), thus

$$\prod_{s=0}^{N_s+1} T_s^{-1} = \prod_{s=1}^{N_s} T_s^{-1} .$$

Explicitly, the transformation (43)-(45) is replaced by (16)-(18) and the action (1) can be expressed as

$$\begin{aligned} \mathcal{A}_f[\eta, \bar{\eta}] &= \bar{\eta}_1 (P_- - rm_q P_+) \eta_1 - \bar{\eta}_1 T_1^{-1} \eta_2 + \sum_{s=2}^{N_s-1} \left\{ \bar{\eta}_s \eta_s - \bar{\eta}_s T_s^{-1} \eta_{s+1} \right\} + \\ & \bar{\eta}_{N_s} \eta_{N_s} - \bar{\eta}_{N_s} T_{N_s}^{-1} (P_+ - rm_q P_-) \eta_1 . \end{aligned}$$

Thus the fermion determinant

$$\det \mathcal{D}(m_q) = \int [d\bar{\psi}][d\psi] e^{-\mathcal{A}_f[\psi, \bar{\psi}]} = \mathcal{J} \int [d\bar{\eta}][d\eta] e^{-\mathcal{A}_f[\eta, \bar{\eta}]}$$

can be evaluated by integrating $\{\eta_s, \bar{\eta}_s\}$ successively from $s = N_s$ to $s = 1$, and the result is given in (19).

As discussed in the Introduction, one cannot identify $(D_c + m_q)$ as the effective 4D lattice Dirac operator for the internal quark fields, since D_c is nonlocal in the limit $N_s \rightarrow \infty$, and $\det(D_c + m_q)$ does not reproduce $\det[\gamma_\mu(\partial_\mu + iA_\mu) + m_q]$ in the continuum limit ($a \rightarrow 0$). The resolution is to introduce a set of pseudofermion fields $\{\phi, \bar{\phi}\}$ similar to the domain-wall fermions but obeying Bose statistics. Explicitly, the action of the pseudofermion fields (with $m_q = 2m_0$) corresponding to the optimal DWF action (34) is

$$\begin{aligned} \mathcal{A}'_{pf} = & \sum_{s,s'=0}^{N_s+1} \sum_{x,x'} \bar{\phi}(x, s) \{ (\omega_s a_5 D_w(x, x') + \delta_{x,x'}) \delta_{s,s'} \\ & - (\delta_{x,x'} - \omega_s a_5 D_w(x, x')) (P_- \delta_{s',s+1} + P_+ \delta_{s',s-1}) \} \phi(x', s') \end{aligned} \quad (54)$$

with boundary conditions

$$\begin{aligned} P_+ \phi(x, -1) &= -P_+ \phi(x, N_s + 1) , \\ P_- \phi(x, N_s + 2) &= -P_- \phi(x, 0) , \end{aligned}$$

where $\omega_0 = \omega_{N_s+1} = 0$, and other weights are given by (7) as before. Thus the integrations over the pseudofermion fields give

$$\int [d\bar{\phi}][d\phi] e^{-\mathcal{A}'_{pf}} = \pi^{N_s+2} K^{-1} \det(1 + rD_c)^{-1} \quad (55)$$

Substituting (51) and (55) into (33), we obtain

$$e^{W[J, \bar{J}]} = \frac{\int [dU] e^{-\mathcal{A}_g} \det[(D_c + m_q)(1 + rD_c)^{-1}] \exp \left\{ \bar{J} r^{-1} (D_c + m_q)^{-1} J \right\}}{\int [dU] e^{-\mathcal{A}_g} \det[(D_c + m_q)(1 + rD_c)^{-1}]} \quad (56)$$

This is one of the main results of this paper.

Now it is clear that the most crucial quantity in (56) is D_c which becomes chirally symmetric in the limit $N_s \rightarrow \infty$. At any finite N_s , D_c (21) preserves the chiral symmetry optimally since S_{opt} (22) is equal to the optimal rational polynomial of the sign function (9). Moreover, $\omega_s a_5$ (7) is independent of a_5 , since it only depends on N_s , λ_{min} and $\lambda_{max}^2/\lambda_{min}^2$. It follows that T_s , S_{opt} , and D_c are all independent of a_5 . Therefore the generating functional $Z[J, \bar{J}]$ for the optimal domain-wall fermion action (34) is a_5 -invariant, preserves the chiral symmetry optimally for any finite N_s . Further, it also manifests the continuum discrete symmetries, since the quark propagator $(D_c + m_q)^{-1}$ (coupling to $\bar{J}$ and J) transforms like $[\gamma_\mu(\partial_\mu + iA_\mu) + m_q]^{-1}$.

From (56), it is clear that the effective 4D lattice Dirac operator for the internal quark fields (dressed with pseudofermions) is

$$D(m_q) = (D_c + m_q)(1 + rD_c)^{-1} = m_q + (m_0 - m_q/2)(1 + \gamma_5 H_w R_Z(H_w)) \quad (57)$$

which is exactly equal to that (25) of the original action (1).

Now any connected n -point Green's function can be obtained by differentiating $W[J, \bar{J}]$ with respect to J and $\bar{J}$ successively. In particular, the quark propagator is

$$\begin{aligned} r\langle q(x)\bar{q}(y)\rangle &= -r \frac{\delta^2 W[J, \bar{J}]}{\delta \bar{J}(x) \delta J(y)} \Big|_{J=\bar{J}=0} \\ &= \frac{\int [dU] e^{-\mathcal{A}_g} \det[D(m_q)] (D_c + m_q)^{-1}(x, y)}{\int [dU] e^{-\mathcal{A}_g} \det[D(m_q)]} \end{aligned} \quad (58)$$

which, in a background gauge field, becomes

$$r\langle q(x)\bar{q}(y)\rangle = (D_c + m_q)^{-1}(x, y) , \quad (59)$$

and it goes to the Dirac propagator $[\gamma_\mu(\partial_\mu + iA_\mu) + m_q]^{-1}(x, y)$ in the continuum limit.

Now it is straightforward to generalize the optimal DWF action (34) to lattice QCD with n_f quark flavors, with quark fields defined as

$$q_f(x) = \sqrt{r}[P_- \psi_f(x, 0) + P_+ \psi_f(x, N_s + 1)] , \quad f = 1, \dots, n_f, \quad (60)$$

$$\bar{q}_f(x) = \sqrt{r}[\bar{\psi}_f(x, 0)P_+ + \bar{\psi}_f(x, N_s + 1)P_-] , \quad (61)$$

where the normalization factor $\sqrt{r} = 1/\sqrt{2m_0}$ has been included. Then the generating functional (56) can be rewritten as

$$Z[\{J_f, \bar{J}_f\}] = \frac{\int [dU] e^{-\mathcal{A}_g} \prod_{f=1}^{n_f} \det[D(m_f)] \exp \left\{ \sum_{f=1}^{n_f} \bar{J}_f (D_c + m_f)^{-1} J_f \right\}}{\int [dU] e^{-\mathcal{A}_g} \prod_{f=1}^{n_f} \det[D(m_f)]} \quad (62)$$

where

$$D(m_f) = (D_c + m_f)(1 + rD_c)^{-1} = m_f + (1 - rm_f)D$$

Evidently, the generating functional (62) ensures that any bilinear (and higher-order) quark operator manifests the continuum discrete symmetries, since the quark fields (60)-(61), both can be decomposed into left and right handed components with the usual chiral projectors, i.e.

$$q_f^L(x) = P_- q_f(x), \quad q_f^R(x) = P_+ q_f(x), \quad (63)$$

$$\bar{q}_f^L(x) = \bar{q}_f(x)P_+, \quad \bar{q}_f^R(x) = \bar{q}_f(x)P_-. \quad (64)$$

Thus the V-A structure of the left-handed quark current

$$L_\mu^{f_1, f_2}(z) = \bar{q}_{f_1}(z) \gamma_\mu (1 - \gamma_5) q_{f_2}(z) \quad (65)$$

in the effective weak Hamiltonian is preserved exactly. This is crucial for determining the indirect CP violation in the $K \rightarrow \pi\pi$ decay through the $\bar{K}^0$ - K^0 mixing. The weak mixing matrix element

$$\langle \bar{K}^0 | \bar{s} \gamma_\mu (1 - \gamma_5) d \bar{s} \gamma_\mu (1 - \gamma_5) d | K^0 \rangle \quad (66)$$

(where s and d denote the strange and down quark fields respectively) can be extracted by evaluating the connected 3-point function (for all $\vec{x}$ and $\vec{y}$ and a range of x_4 and y_4)

$$\begin{aligned} & \langle \bar{d}(x) \gamma_4 \gamma_5 s(x) L_\mu^{s,d}(0) L_\mu^{s,d}(0) \bar{d}(y) \gamma_4 \gamma_5 s(y) \rangle^{\text{connected}} \\ &= \frac{\delta}{\delta J_d(x)} \gamma_4 \gamma_5 \frac{\delta}{\delta \bar{J}_s(x)} \frac{\delta}{\delta J_s(0)} \gamma_\mu (1 - \gamma_5) \frac{\delta}{\delta \bar{J}_d(0)} \times \\ & \quad \frac{\delta}{\delta J_s(0)} \gamma_\mu (1 - \gamma_5) \frac{\delta}{\delta \bar{J}_d(0)} \frac{\delta}{\delta J_d(y)} \gamma_4 \gamma_5 \frac{\delta}{\delta \bar{J}_s(y)} Z[\{J_f, \bar{J}_f\}] \Big|_{J=\bar{J}=0} \\ & \quad - \text{disconnected diagrams} \\ &= \frac{1}{\mathcal{Z}} \int [dU] e^{-\mathcal{A}_g} \prod_f \det[D(m_f)] \text{tr}[(D_c + m_d)_{0,x}^{-1} \gamma_4 \gamma_5 (D_c + m_s)_{x,0}^{-1} \gamma_\mu (1 - \gamma_5)] \times \\ & \quad \text{tr}[(D_c + m_s)_{y,0}^{-1} \gamma_\mu (1 - \gamma_5) (D_c + m_d)_{0,y}^{-1} \gamma_4 \gamma_5] , \end{aligned}$$

and the 2-point functions

$$\begin{aligned} & \langle \bar{d}(x) \gamma_4 \gamma_5 s(x) \bar{s}(0) \gamma_4 \gamma_5 d(0) \rangle \\ &= \frac{1}{\mathcal{Z}} \int [dU] e^{-\mathcal{A}_g} \prod_f \det[D(m_f)] \text{tr}[(D_c + m_d)_{0,x}^{-1} \gamma_4 \gamma_5 (D_c + m_s)_{x,0}^{-1} \gamma_4 \gamma_5] , \\ & \quad \langle \bar{s}(0) \gamma_4 \gamma_5 d(0) \bar{d}(y) \gamma_4 \gamma_5 s(y) \rangle \\ &= \frac{1}{\mathcal{Z}} \int [dU] e^{-\mathcal{A}_g} \prod_f \det[D(m_f)] \text{tr}[(D_c + m_s)_{y,0}^{-1} \gamma_4 \gamma_5 (D_c + m_d)_{0,y}^{-1} \gamma_4 \gamma_5] , \end{aligned}$$

where

$$\mathcal{Z} = \int [dU] e^{-\mathcal{A}_g} \prod_f \det[D(m_f)] .$$

It is clear that there is *no* right-handed currents entering the weak mixing matrix element (66), and the $SU(2)_L$ gauge symmetry of electroweak interactions is preserved exactly. Had one *not* introduced the boundary layers at $s = 0$ and $s = N_s + 1$, then one might *not* have found any q and $\bar{q}$ such that Eq. (59) is satisfied and the usual chiral projectors $(1 \pm \gamma_5)/2$ can work for both q and $\bar{q}$. Consequently its left-handed quark current may not manifest the V-A structure, and the $SU(2)_L$ gauge symmetry may be broken by the order $O(a)$ effect. Obviously, such left-handed quark current is not ideal for determining the $\bar{K}^0$ - K^0 mixing.

In practice, the valence quark propagator $(D_c + m_q)^{-1}$ in a background gauge field can be computed via the 5-dimensional optimal DWF operator $\mathcal{D}'$ (34). Using (59) and the definition of quark fields (37)-(38), one immediately obtains

$$\begin{aligned}
& r^{-1}(D_c + m_q)^{-1}(x, y) \\
&= \langle q(x) \bar{q}(y) \rangle \\
&= \sum_{s, s'} (P_- \delta_{s,0} + P_+ \delta_{s, N_s+1}) \frac{\int [d\bar{\psi}] [d\psi] \psi_s(x) \bar{\psi}_{s'}(y) e^{-\mathcal{A}'_f}}{\int [d\bar{\psi}] [d\psi] e^{-\mathcal{A}'_f}} (P_+ \delta_{s',0} + P_- \delta_{s', N_s+1}) \\
&= \sum_{s, s'} (P_- \delta_{s,0} + P_+ \delta_{s, N_s+1}) \mathcal{D}'^{-1}(x, s; y, s') (P_+ \delta_{s',0} + P_- \delta_{s', N_s+1}) , \tag{67}
\end{aligned}$$

where $\mathcal{D}'^{-1}(x, s; y, s')$ can be obtained by solving the linear system

$$\mathcal{D}'(x, s; y, s') Y = \mathbb{1} . \tag{68}$$

Alternatively, one can also obtain the valence quark propagator

$$(D_c + m_q)^{-1} = (1 - r m_q)^{-1} [D^{-1}(m_q) - r] , \tag{69}$$

by computing $D^{-1}(m_q)$ via the original optimal DWF operator (1), i.e., solving the linear system

$$\mathcal{D}(m_q) Y = \mathcal{D}(2m_0) B^{-1} \mathbb{1} , \tag{70}$$

where B and B^{-1} are defined in (114) and (115) respectively. Then the solution Y gives

$$D^{-1}(m_q)(x, x') = r(BY)(x, 1; x', 1) , \tag{71}$$

The derivation of (70) is given in the Appendix.

Note that we can also compute the valence quark propagator (69) by solving the linear system

$$D(m_q) Y = \left[m_q + (m_0 - m_q/2) \left(1 + \gamma_5 H_w R_Z(H_w^2) \right) \right] Y = \mathbb{1} \tag{72}$$

where $D(m_q)$ is the effective 4D lattice Dirac operator (57), and $R_Z(H_w^2)$ is the Zolotarev rational polynomial as defined in (9)-(11).

Since solving any one of above three linear systems (72), (68) or (70) yields the same quark propagator, in principle, it does *not* matter which linear system one chooses. However, in practice, one should pick the most efficient scheme for one's computational system (hardware and software).

2.2 Conventional domain-wall fermions

Shamir's domain-wall fermion action [4] can be written as

$$\begin{aligned}
\mathcal{A}_{fs} &= \sum_{s,s'=1}^{N_s} \sum_{x,x'} \bar{\psi}(x,s) [(a_5 D_w(x,x') + \delta_{x,x'}) \delta_{s,s'} \\
&\quad - (P_- \delta_{s',s+1} + P_+ \delta_{s',s-1}) \delta_{x,x'}] \psi(x',s') \\
&\equiv \sum_{s,s'=1}^{N_s} \sum_{x,x'} \bar{\psi}(x,s) \mathcal{D}_{fs}(x,s; x',s') \psi(x',s')
\end{aligned} \tag{73}$$

with boundary conditions

$$P_+ \psi(x, 0) = -rm_q P_+ \psi(x, N_s), \quad r = \frac{1}{2m_0} \tag{74}$$

$$P_- \psi(x, N_s + 1) = -rm_q P_- \psi(x, 1), \tag{75}$$

where D_w is defined in (4), and m_q is the bare quark mass. The quark fields are constructed from the boundary modes:

$$q(x) = P_- \psi(x, 1) + P_+ \psi(x, N_s), \tag{76}$$

$$\bar{q}(x) = \bar{\psi}(x, 1) P_+ + \bar{\psi}(x, N_s) P_- . \tag{77}$$

The fermion determinant and the quark propagator (in gauge background) have been derived in Refs. [15]-[17]. Now we derive the generating functional for n -point Green's function of the quarks fields, according to the formula (33). The derivation is similar to that of the optimal DWF action in the last subsection.

Using $\gamma_5 P_{\pm} = \pm P_{\pm}$, $P_+ P_- = 0$, $P_+ + P_- = 1$, and $H_w = \gamma_5 D_w$, we can rewrite (73) as

$$\begin{aligned}
\mathcal{A}_{fs} &= -(rm_q + 1) \bar{\psi}_1 \gamma_5 (a_5 H_w P_- - 1) P_+ \psi_{N_s} \\
&\quad + \bar{\psi}_1 \gamma_5 (a_5 H_w P_- - 1) (P_- \psi_1 + P_+ \psi_{N_s}) + \bar{\psi}_1 \gamma_5 (a_5 H_w P_+ + 1) (P_- \psi_2 + P_+ \psi_1) \\
&\quad + \sum_{s=2}^{N_s-1} \left\{ \bar{\psi}_s \gamma_5 (a_5 H_w P_- - 1) (P_- \psi_s + P_+ \psi_{s-1}) + \bar{\psi}_s \gamma_5 (a_5 H_w P_+ + 1) (P_- \psi_{s+1} + P_+ \psi_s) \right\} \\
&\quad + \bar{\psi}_{N_s} \gamma_5 (a_5 H_w P_- - 1) (P_- \psi_{N_s} + P_+ \psi_{N_s-1}) + \bar{\psi}_{N_s} \gamma_5 (a_5 H_w P_+ + 1) (P_- \psi_1 + P_+ \psi_{N_s}) \\
&\quad - (rm_q + 1) \bar{\psi}_{N_s} \gamma_5 (a_5 H_w P_+ + 1) P_- \psi_1
\end{aligned} \tag{78}$$

where all indices are suppressed except the flavor index in the fifth dimension.

Next we define

$$\begin{aligned}
\eta_1 &= P_- \psi_1 + P_+ \psi_{N_s}, \quad \bar{\eta}_1 = \bar{\psi}_1 \gamma_5 (a_5 H_w P_- - 1), \\
\eta_s &= P_- \psi_s + P_+ \psi_{s-1}, \quad \bar{\eta}_s = \bar{\psi}_s \gamma_5 (a_5 H_w P_- - 1), \quad s = 2, \dots, N_s - 1 \\
\eta_{N_s} &= P_- \psi_{N_s} + P_+ \psi_{N_s-1}, \quad \bar{\eta}_{N_s} = \bar{\psi}_{N_s} \gamma_5 (a_5 H_w P_- - 1),
\end{aligned}$$

and the inverse transform

$$\begin{aligned}\psi_1 &= P_- \eta_1 + P_+ \eta_2, \quad \bar{\psi}_1 = \bar{\eta}_1 (a_5 H_w P_- - 1)^{-1} \gamma_5, \\ \psi_s &= P_- \eta_s + P_+ \eta_{s+1}, \quad \bar{\psi}_s = \bar{\eta}_s (a_5 H_w P_- - 1)^{-1} \gamma_5, \quad s = 2, \dots, N_s - 1 \\ \psi_{N_s} &= P_- \eta_{N_s} + P_+ \eta_1, \quad \bar{\psi}_{N_s} = \bar{\eta}_{N_s} (a_5 H_w P_- - 1)^{-1} \gamma_5.\end{aligned}$$

Then the action (78) can be expressed in terms of $\{\eta, \bar{\eta}\}$ fields

$$\begin{aligned}\mathcal{A}_{fs} &= \bar{\eta}_1 (P_- - rm_q P_+) \eta_1 - \bar{\eta}_1 T^{-1} \eta_2 + \sum_{s=2}^{N_s-1} \left\{ \bar{\eta}_s \eta_s - \bar{\eta}_s T^{-1} \eta_{s+1} \right\} \\ &\quad + \bar{\eta}_{N_s} \eta_{N_s} - \bar{\eta}_{N_s} T^{-1} (P_+ - rm_q P_-) \eta_1\end{aligned}\tag{79}$$

where

$$T = (1 + a_5 H_w P_+)^{-1} (1 - a_5 H_w P_-) = \frac{1 - a_5 H}{1 + a_5 H},\tag{80}$$

$$H \equiv (2 + a_5 H_w \gamma_5)^{-1} H_w = H_w (2 + \gamma_5 a_5 H_w)^{-1} = \gamma_5 \frac{D_w}{2 + a_5 D_w}.\tag{81}$$

At this point, it is instructive to compare the transfer matrix T (80) with its counterpart T_s (23) in the optimal DWF. As it will become clear later, one of the difficulties in the conventional DWF is due to the fact that the denominator of H (81) is $(2 + a_5 D_w)$ rather than a constant.

Now the fermionic integral in the numerator of (33) can be written as

$$\int [d\bar{\psi}] [d\psi] e^{-\mathcal{A}_{fs} + \bar{J}q + \bar{q}J} = \tau \int [d\bar{\eta}] [d\eta] e^{-\mathcal{A}_{fs} + \bar{J}\eta_1 - \bar{\eta}_1 P_+ J + \bar{\eta}_{N_s} T^{-1} P_- J}\tag{82}$$

where

$$\begin{aligned}\bar{J}q + \bar{q}J &= \bar{J}(P_- \psi_1 + P_+ \psi_{N_s}) + (\bar{\psi}_1 P_+ + \bar{\psi}_{N_s} P_-) J \\ &= \bar{J}\eta_1 + \bar{\eta}_1 (a_5 H_w P_- - 1)^{-1} P_+ J - \bar{\eta}_{N_s} (a_5 H_w P_- - 1)^{-1} P_- J \\ &= \bar{J}\eta_1 - \bar{\eta}_1 P_+ J + \bar{\eta}_{N_s} T^{-1} P_- J,\end{aligned}$$

and τ is the Jacobian of the transformation,

$$\tau = \det [\gamma_5 (a_5 H_w P_- - 1)]^{N_s}.$$

Again using the Grassman integral formula (50), one can easily evaluate the Grassman integrals in (82), by integrating $(\eta_s, \bar{\eta}_s)$ successively from $s = N_s$ to $s = 1$, similar to the steps in the last subsection. The final result is

$$\begin{aligned}&\tau \det [(P_- - rm_q P_+) - T^{-N_s} (P_+ - rm_q P_-)] \times \\ &\exp \left\{ \bar{J} [(-P_+ + T^{-N_s} P_-)^{-1} (P_- - T^{-N_s} P_+) + rm_q]^{-1} J \right\} \\ &= K_f \det [r(D_c^{(s)} + m_q)] \exp \left\{ \bar{J} r^{-1} (D_c^{(s)} + m_q)^{-1} J \right\}\end{aligned}\tag{83}$$

where

$$K_f = \det[\gamma_5(a_5 H_w P_- - 1)]^{N_s} \det(-P_+ + T^{-N_s} P_-) \quad (84)$$

$$rD_c^{(s)} = (-P_+ + T^{-N_s} P_-)^{-1} (P_- - T^{-N_s} P_+) = \frac{1 + \gamma_5 S(H)}{1 - \gamma_5 S(H)} \quad (85)$$

$$S(H) = \frac{1 - T^{N_s}}{1 + T^{N_s}} = \frac{(1 + a_5 H)^{N_s} - (1 - a_5 H)^{N_s}}{(1 + a_5 H)^{N_s} + (1 - a_5 H)^{N_s}}. \quad (86)$$

Again, since $D_c^{(s)}$ is nonlocal in the limit $N_s \rightarrow \infty$, one has to introduce pseudofermion fields (similar to the domain-wall fermions but obeying Bose statistics) with action (73) ($m_q = 2m_0$) such that the effective 4D lattice Dirac operator for the internal fermion loops is exponentially-local for smooth gauge backgrounds, and it goes to $[\gamma_\mu(\partial_\mu + iA_\mu) + m_q]$ in the continuum limit $a, a_5 \rightarrow 0$. Then integrating the pseudofermion fields gives

$$\int [d\bar{\phi}][d\phi] e^{-\mathcal{A}_{pfs}} = \pi^{N_s} K_f^{-1} \det(1 + rD_c^{(s)})^{-1}. \quad (87)$$

Substituting (83) and (87) into (33), we obtain

$$e^{W[J, \bar{J}]} = \frac{\int [dU] e^{-\mathcal{A}_g} \det[(D_c^{(s)} + m_q)(1 + rD_c^{(s)})^{-1}] \exp\{\bar{J} r^{-1} (D_c^{(s)} + m_q)^{-1} J\}}{\int [dU] e^{-\mathcal{A}_g} \det[(D_c^{(s)} + m_q)(1 + rD_c^{(s)})^{-1}]} \quad (88)$$

which is exactly in the same form of (56) except replacing D_c with $D_c^{(s)}$ [i.e., replacing S_{opt} (22) with S (86)]. However, such a difference has significant consequences, as we will demonstrate in the next section.

In the limit $N_s \rightarrow \infty$, $S(H)$ (86) becomes the sign function of H ,

$$\lim_{N_s \rightarrow \infty} S(H) = \frac{H}{\sqrt{H^2}},$$

thus $D_c^{(s)}$ (85) is chirally symmetric. However, *it still depends on a_5 through H* (81). In other words, the lattice spacing a_5 in the conventional DWF could be a relevant parameter² which must be taken to zero before its effective 4D lattice Dirac operator

$$D^{(s)}(m_q) = (D_c^{(s)} + m_q)(1 + rD_c^{(s)})^{-1} \quad (89)$$

can behave properly.

However, taking the limit $a_5 \rightarrow 0$ is rather tedious in practice. Therefore, it is vital to construct a domain-wall fermion action such that its generating

²As demonstrated by the tests in Section 3, the fermion determinant has a strong dependence on a_5 for 2D topologically nontrivial gauge configurations.

functional $Z[J, \bar{J}]$ (thus any n -point Green's function) is independent of a_5 , for any a and N_s .

This objective can be partially fulfilled if H (81) is replaced by $H_w = \gamma_5 D_w$. Then, in the limit $N_s \rightarrow \infty$, $S(H_w)$ (86) becomes independent of a_5 ,

$$\lim_{N_s \rightarrow \infty} S(H_w) = \frac{a_5 H_w}{\sqrt{a_5^2 H_w^2}} = \frac{H_w}{\sqrt{H_w^2}} .$$

However, at finite N_s , $S(H_w)$ still depends on a_5 .

Aiming at modifying the action (73) such that H is replaced by H_w in the transfer matrix T , one immediately observes that if all factors $(a_5 H_w P_{\pm} \pm 1)$ in the action (78) are replaced by $(a_5 H_w \pm 1)$, then T becomes

$$T' = \frac{1 - a_5 H_w}{1 + a_5 H_w} . \quad (90)$$

This amounts to adding the following terms to the DWF action (78),

$$\begin{aligned} \delta \mathcal{A}_{f_s} = & -(rm_q + 1) \bar{\psi}_1 \gamma_5 a_5 H_w P_+ \psi_{N_s} + \\ & + \bar{\psi}_1 \gamma_5 a_5 H_w P_+ (P_- \psi_1 + P_+ \psi_{N_s}) + \bar{\psi}_1 \gamma_5 a_5 H_w P_- (P_- \psi_2 + P_+ \psi_1) \\ & + \sum_{s=2}^{N_s-1} \left\{ \bar{\psi}_s \gamma_5 a_5 H_w P_+ (P_- \psi_s + P_+ \psi_{s-1}) + \bar{\psi}_s \gamma_5 a_5 H_w P_- (P_- \psi_{s+1} + P_+ \psi_s) \right\} \\ & + \bar{\psi}_{N_s} \gamma_5 a_5 H_w P_+ (P_- \psi_{N_s} + P_+ \psi_{N_s-1}) + \bar{\psi}_{N_s} \gamma_5 a_5 H_w P_- (P_- \psi_1 + P_+ \psi_{N_s}) \\ & - (rm_q + 1) \bar{\psi}_{N_s} \gamma_5 a_5 H_w P_- \psi_1 , \end{aligned}$$

which can be simplified to

$$\begin{aligned} \delta \mathcal{A}_{f_s} = & -rm_q \bar{\psi}_1 \gamma_5 a_5 H_w P_+ \psi_{N_s} + \bar{\psi}_1 \gamma_5 a_5 H_w P_- \psi_2 \\ & + \sum_{s=2}^{N_s-1} \left\{ \bar{\psi}_s \gamma_5 a_5 H_w P_+ \psi_{s-1} + \bar{\psi}_s \gamma_5 a_5 H_w P_- \psi_{s+1} \right\} \\ & + \bar{\psi}_{N_s} \gamma_5 a_5 H_w P_+ \psi_{N_s-1} - rm_q \bar{\psi}_{N_s} \gamma_5 a_5 H_w P_- \psi_1 . \end{aligned} \quad (91)$$

Then adding (91) to (78), one obtains

$$\begin{aligned} \mathcal{A}'_{f_s} = & \sum_{s,s'=1}^{N_s} \sum_{x,x'} \bar{\psi}(x, s) \{ (a_5 D_w(x, x') + \delta_{x,x'}) \delta_{s,s'} \\ & - (\delta_{x,x'} - a_5 D_w(x, x')) (P_+ \delta_{s',s-1} + P_- \delta_{s',s+1}) \} \psi(x', s') , \end{aligned} \quad (92)$$

which agrees with Borici's domain-wall action [18]. The effective 4D lattice Dirac operator of (92) is the same as (89) except (the argument of $D_c^{(s)}$) $S(H)$ (86) is replaced by $S(H_w)$. Thus, at $a_5 = 1$, it (times a factor $r = 1/2m_0$) is

exactly equal to the overlap Dirac operator with $(H_w^2)^{-1/2}$ approximated by the polar approximation, since

$$S(H_w) = \begin{cases} a_5 H_w \left(\frac{1}{N_s} + \frac{2}{N_s} \sum_{l=1}^n \frac{b_l}{a_5^2 H_w^2 + d_l} \right) , & N_s = 2n + 1 \text{ (odd)} \\ a_5 H_w \frac{2}{N_s} \sum_{l=1}^n \frac{b_l}{a_5^2 H_w^2 + d_l} , & N_s = 2n \text{ (even)} \end{cases}$$

where

$$b_l = \sec^2 \left[\frac{\pi}{N_s} \left(l - \frac{1}{2} \right) \right] , \quad d_l = \tan^2 \left[\frac{\pi}{N_s} \left(l - \frac{1}{2} \right) \right] .$$

Obviously, for any finite N_s , $S(H_w)$ depends on a_5 . Also, it does not preserve chiral symmetry optimally, since $S(H_w)$ is not the optimal rational approximation of $\text{sgn}(H_w)$, as shown in Ref. [5].

A theoretical inconsistency of the Borici DWF action (92) is that the quark propagator $(D'_c + m_q)^{-1}$ *cannot* be expressed in terms of the correlation function of the quark fields constructed from the boundary modes (76)-(77), i.e.,

$$r \langle q(x) \bar{q}(y) \rangle \neq (D'_c + m_q)_{x,y}^{-1} , \quad r D'_c = \frac{1 + \gamma_5 S(H_w)}{1 - \gamma_5 S(H_w)} ,$$

in any background gauge field. Thus the generating functional (with localized quark sources $J(x)$ and $\bar{J}(x)$) for n -point Green's function of q and $\bar{q}$ is not well-defined, and *cannot* be derived.

In this paper, a solution to this problem is presented in the last subsection. Explicitly, one can modify the action (92) by adding two boundary layers with identity transfer matrix (i.e., without $a_5 D_w$ couplings) at $s = 0$ and $s = N_s + 1$, with boundary conditions (35)-(36), and defining the quark fields from the new boundary modes (37)-(38), similar to the new optimal DWF (34). Then the generating functional for n -point Green's function of q and $\bar{q}$ can be derived in the same way as that presented in the last subsection, and the result is in the same form of (56) except replacing each $T_s(s = 1, \dots, N_s)$ (23) with T' (90).

3 Tests

In this section, we compute the index, the axial anomaly and the fermion determinant of the effective lattice Dirac operator of the optimal DWF (34)³, and the conventional DWF (73) respectively, in two-dimensional background $U(1)$ gauge fields. Our notations for the two-dimensional background gauge field are the same as those in Eqs. (7)-(11) of Ref. [20].

³Note that both (1) and (34) have the same effective lattice Dirac operator (25), as shown in Section 2.

The effective lattice Dirac operator of the optimal DWF can be computed from the 5-dimensional DWF operator (34). Setting $m_q = 2m_0$ ($rm_q = 1$) in (67), one has

$$(1 + rD_c)^{-1}(x, y) = \sum_{s, s'} (P_- \delta_{s,0} + P_+ \delta_{s, N_s+1}) \{ \mathcal{D}'^{-1}(2m_0) \} (x, s; y, s') (P_+ \delta_{s',0} + P_- \delta_{s', N_s+1}).$$

Then the massless lattice Dirac operator can be obtained by a subtraction

$$D = D_c(1 + rD_c)^{-1} = 1 - (1 + rD_c)^{-1}, \quad (93)$$

and the massive one by

$$D(m_q) = (D_c + m_q)(1 + rD_c)^{-1} = 1 + (m_q - 1)(1 + rD_c)^{-1}. \quad (94)$$

Similarly, we can obtain the effective lattice Dirac operator of the conventional DWF (73), with the boundary modes at $s = 1$ and N_s .

In the following tests, we fix $m_0 = 1$ ($r = 1/2$) in D_w (4), and $m_q = 0$ (except for the fermion determinants in Table 2 with $m_q = 0.05$). With N_s in the range $10 \leq N_s \leq 32$, we first check that the eigenvalues of the effective lattice Dirac operators fall on a circle of radius one ($1/2r = 1$) centered at one. For nontrivial gauge backgrounds, the real eigenmodes (zero and two) have definite (+1 or -1) chirality and satisfy the chirality sum rule [20]

$$n_+ + N_+ = n_- + N_- , \quad (95)$$

where $n_{\pm}(N_{\pm})$ stand for the numbers of zero (+2) eigenmodes with ± 1 chirality respectively. These real eigenmodes are topological, i.e., robust under local fluctuations of the gauge fields. For any (trivial or nontrivial) gauge background, the complex eigenmodes must come in conjugate pairs, each with zero chirality. The Atiyah-Singer index theorem ($n_+ - n_- = Q$) holds for any gauge configuration (with topological charge Q) satisfying the topological bound [22]

$$\forall_x |\bar{\rho}(x)| < 0.28 \quad (96)$$

where $\bar{\rho}(x)$ is the topological charge (density) inside a unit square (of area a^2) centered at x ,

$$\bar{\rho}(x) = \frac{1}{a^2} \int_{x_1-a/2}^{x_1+a/2} dy_1 \int_{x_2-a/2}^{x_2+a/2} dy_2 \frac{1}{2\pi} F_{12}(y). \quad (97)$$

Thus, we conclude that the effective lattice Dirac operators of the optimal DWF and the conventional DWF both are topologically-proper for sufficiently large N_s .

3.1 Axial anomaly

Since the effective lattice Dirac operator D of domain-wall fermions ($N_s \rightarrow \infty$) satisfies the Ginsparg-Wilson relation (29), its axial anomaly (for any N_s) can be defined as [19]

$$\mathcal{A}_L(x) = \text{tr}[\gamma_5(1 - rD(x, x))] = -\frac{1}{2}\text{tr}[\mathcal{S}(x, x)] , \quad (98)$$

where the trace runs over Dirac and color indices. Here the operator $\mathcal{S}$ becomes S_{opt} (22) for the optimal DWF, and S (86) for the conventional DWF.

For sufficiently large N_s and for gauge configurations satisfying the topological bound (96), I checked that the sum of the axial anomaly over all sites is equal to the index of D and the topological charge Q

$$\sum_x \mathcal{A}_L(x) = n_+ - n_- = Q , \quad (99)$$

for both the optimal DWF and the conventional DWF. However, this does not necessarily imply that $\mathcal{A}_L(x)$ is equal to the topological charge density at each site.

In the following, I compute the axial anomaly $\mathcal{A}_L(x)$ for the overlap Dirac operator, the optimal DWF, and the conventional DWF respectively, and compare them with the topological charge density $\bar{\rho}(x)$ (97) of the gauge background on the torus. The deviation of the axial anomaly can be measured in terms of

$$\delta = \frac{1}{N} \sum_x |\mathcal{A}_L(x) - a^2 \bar{\rho}(x)| \quad (100)$$

where $\bar{\rho}(x)$ is the topological charge density inside a unit square (of area a^2) centered at x .

In a nontrivial $U(1)$ gauge background with parameters $Q = 1$, $h_1 = 0.1$, $h_2 = 0.2$, $A_1^{(0)} = 0.3$, $A_2^{(0)} = 0.4$ and $n_1 = n_2 = 1$ (as defined in Eqs. (7) and (8) in Ref. [20]) on the 8×8 lattice, the axial anomaly $\mathcal{A}_L(x)$ and its deviation δ are computed for the overlap, the optimal DWF and the conventional DWF respectively. Here the 2D overlap Dirac operator is computed exactly via diagonalization.

The results are :

$$\delta = \begin{cases} 4.75 \times 10^{-3}, & \text{overlap} , \\ 4.75 \times 10^{-3}, & \text{optimal DWF } (N_s = 16) , \\ 5.44 \times 10^{-3}, & \text{conventional DWF } (N_s = 16, 32) . \end{cases} \quad (101)$$

The slightly larger deviation of the axial anomaly of the conventional DWF indicates that its effective lattice Dirac operator is less localized than that of the optimal DWF. Note that there is no improvements by increasing $N_s = 16$ to $N_s = 32$.

For all background gauge configurations we have tested, the optimal DWF always gives anomaly deviation (δ) smaller than that of the conventional DWF.

3.2 Fermion determinant

In quantum field theory, the fermion determinant $\det(D)$ is proportional to the exponentiation of the one-loop effective action which is the summation of any number of external sources interacting with one internal fermion loop. Thus it is one of the most crucial quantities to be examined in any lattice fermion formulations. Even though a (massless) lattice Dirac operator is topologically-proper (i.e., with the correct index and axial anomaly), it does not necessarily imply that its non-zero complex eigenvalues behave properly. This can be tested by computing its fermion determinant

$$\det(D) = (1 + e^{i\pi})^{(n_+ + n_-)} \det'(D) \quad (102)$$

where $\det'(D)$ is equal to the product of all non-zero eigenvalues. Since the effective lattice Dirac operators of the optimal DWF and the conventional DWF are γ_5 -hermitian, the eigenvalues of D are either real or come in complex conjugate pairs, and $\det'(D)$ must be real and positive. For $Q = 0$, we have $n_+ + n_- = 0$ and $\det(D) = \det'(D)$. For $Q \neq 0$, we have $n_+ + n_- \neq 0$ and $\det(D) = 0$, but $\det'(D)$ still provides important information about the spectrum. In continuum, exact solution of fermion determinant in the general background $U(1)$ gauge fields on a torus ($L_1 \times L_2$) was obtained in Ref. [21].

With nontrivial gauge backgrounds, we compute $\det'(D)$ for the overlap Dirac operator, the optimal DWF, and the conventional DWF respectively, and compare them with the exact solution in continuum. For simplicity, we turn off the harmonic part ($h_1 = h_2 = 0$) and the local sinusoidal fluctuations ($A_1^{(0)} = A_2^{(0)} = 0$), and examine the change of $\det'(D)$ with respect to the topological charge Q . For these gauge configurations, the exact solution [21] is

$$\det'[D(Q)] = \mathcal{N} \left(\frac{L_1 L_2}{2|Q|} \right)^{|Q|/2} \quad (103)$$

where the normalization constant $\mathcal{N}$ is fixed by

$$\mathcal{N} = \sqrt{\frac{2}{L_1 L_2}}$$

such that $\det'[D(1)] = 1$.

In Table 1, the fermion determinants $\det'(D)$ on the 8×8 lattice are listed. It is clear that the fermion determinant of the optimal DWF ($N_s = 10$) highly agrees with that of the (exact) overlap Dirac operator, and both of them are in good agreement with the exact solution on the 8×8 torus. On the other hand, the fermion determinant of the conventional DWF ($N_s = 10$, $a_5 = 1$) does not agree with the exact solution. Further, the discrepancies do not decrease as N_s is increased. This suggests that the error is not due to finite N_s , but the

	exact	overlap	optimal DWF	conventional DWF		
Q			$N_s = 10$	$N_s = 10$ $a_5 = 1$	$N_s = 32$ $a_5 = 1$	$N_s = 20$ $a_5 = 0.5$
1	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000
2	2.82843	2.79094	2.79094	1.44349	1.44558	2.08267
3	6.15840	5.96482	5.96482	1.63969	1.64629	3.32119
4	11.3137	10.7257	10.7257	1.61127	1.62411	4.46412
5	18.3179	16.9800	16.9800	1.43207	1.45172	5.29955
6	26.8177	24.3215	24.3215	1.18079	1.20677	5.71414
7	36.1083	32.0978	32.0978	0.91741	0.94853	5.69848
8	45.2548	40.5552	40.5551	0.67521	0.71130	5.37711
9	53.2732	45.9271	45.9269	0.47955	0.51653	4.68732
10	59.3164	50.6766	50.6761	0.32497	0.36279	3.91209

Table 1: The fermion determinant versus the topological charge Q . The normalization constant is chosen such that $\det' D = 1$ for $Q = 1$. The results on the 8×8 lattice are listed for the overlap Dirac operator, the effective lattice Dirac operators of the optimal DWF and the conventional DWF respectively. The exact solutions on the 8×8 torus are computed according to Eq. (103).

lattice spacing a_5 which enters the effective lattice Dirac operator (89) through $S(H)$ (86). If this is the case, then decreasing a_5 would improve the results. This is demonstrated in the last column with $a_5 = 0.5$ and $N_s = 20$, which gives results much better than those of the column with $N_s = 32, a_5 = 1$.

In Table 2, the fermion determinants for massive fermion with $m_q = 0.05$ are listed, for the same nontrivial gauge backgrounds in Table 1. The massive overlap Dirac operator is $[m_q + (1 - rm_q)D_o]$, and the effective lattice Dirac operators of the optimal DWF and the conventional DWF are computed via (94). In each case (column), the normalization constant is fixed such that $\det[D(m_q)] = 1$ for $Q = 0$. Clearly, the fermion determinant of the optimal DWF ($N_s = 10$) is in excellent agreement with that of the (exact) overlap, while the conventional DWF disagrees with the overlap and the optimal DWF. Further, the discrepancies do not decrease as N_s is increased. However, if a_5 is reduced from 1 to 0.5, then the results are significantly improved, as shown in the last column. This demonstrates that the massive fermion determinant of the conventional DWF is also sensitive to a_5 , for topologically nontrivial gauge configurations.

On the other hand, for 2D topologically trivial gauge configurations, the fermion determinant of the conventional DWF only has a weak dependence on a_5 . This can be demonstrated as follows. For simplicity, we turn off the local sinusoidal fluctuations ($A_1^{(0)} = A_2^{(0)} = 0$), and examine the change of $\det(D)$ versus the harmonic parts $h_1 = h_2$. In Table 3, the fermion determinants

	overlap	optimal DWF	conventional DWF		
Q		$N_s = 10$	$N_s = 10$ $a_5 = 1$	$N_s = 32$ $a_5 = 1$	$N_s = 20$ $a_5 = 0.5$
0	1.00000	1.00000	1.00000	1.00000	1.00000
1	5.27834	5.27834	10.4991	10.4943	7.00364
2	0.73042	0.73042	0.75612	0.75684	0.72616
3	7.768×10^{-2}	7.768×10^{-2}	4.289×10^{-2}	4.304×10^{-2}	5.782×10^{-2}
4	6.961×10^{-3}	6.961×10^{-3}	2.106×10^{-3}	2.121×10^{-3}	3.893×10^{-3}
5	5.497×10^{-4}	5.497×10^{-4}	9.371×10^{-5}	9.474×10^{-5}	2.332×10^{-4}
6	3.929×10^{-5}	3.929×10^{-5}	3.883×10^{-6}	3.935×10^{-6}	1.289×10^{-5}
7	2.588×10^{-6}	2.588×10^{-6}	1.535×10^{-7}	1.535×10^{-7}	6.830×10^{-7}
8	1.633×10^{-7}	1.633×10^{-7}	5.621×10^{-9}	5.794×10^{-9}	3.407×10^{-8}
9	9.235×10^{-9}	9.235×10^{-9}	2.405×10^{-10}	2.103×10^{-10}	2.156×10^{-9}

Table 2: The fermion determinant $\det[D(m_q)]$ versus the topological charge Q for effective lattice Dirac operator with bare mass $m_q = 0.05$ on the 8×8 lattice. The normalization constant is chosen such that $\det D(m_q) = 1$ for $Q = 0$. The results are listed for the overlap, the optimal DWF ($N_s = 10$), and the conventional DWF with $(N_s = 10, a_5 = 1)$, $(N_s = 32, a_5 = 1)$ and $(N_s = 20, a_5 = 0.5)$ respectively.

versus the harmonic parts ($h_1 = h_2$) on a 8×8 torus (lattice) are listed for the exact solution [21], the overlap, the optimal DWF, and the conventional DWF respectively. Clearly, all results are in very good agreement with each other. Further, the fermion determinant of the conventional DWF is insensitive to a_5 , as comparing the corresponding entries of the last two columns ($N_s = 10, a_5 = 1$) and ($N_s = 20, a_5 = 0.5$).

Even though above tests are performed with 2D abelian gauge backgrounds, one expects that the a_5 effects might also occur in the quark propagator (as well as the fermion determinant), in lattice QCD with the conventional DWF. We are now computing the quark propagator for several different values of a_5 , on the $8^3 \times 24$ lattice, for 100 gauge configurations. Then we will extract the pion mass as well as other observables, and study their dependence on a_5 . These results will be presented elsewhere.

4 Concluding Remarks

To formulate lattice QCD with domain-wall fermions, it seems to be inevitable to introduce parameters N_s and a_5 (in the fifth dimension) to the theory. Presumably, only in the limit $N_s \rightarrow \infty$ and $a_5 \rightarrow 0$, the effective 4D theory agrees with the target theory (QCD), up to the finite volume and discretization

	exact	overlap	optimal DWF	conventional DWF	
$h_1 = h_2$			$N_s = 10$	$N_s = 10$ $a_5 = 1$	$N_s = 20$ $a_5 = 0.5$
0.0	0.00000	0.00000	0.00000	0.00000	0.00000
0.1	1.00000	1.00000	1.00000	1.00000	1.00000
0.2	3.34414	3.40518	3.40518	3.39253	3.40883
0.3	5.71957	5.97173	5.97137	5.92350	5.98565
0.4	7.24506	7.73701	7.73701	7.64986	7.76372
0.5	7.73926	8.34167	8.34166	8.23777	8.37383
0.6	7.24506	7.73701	7.73701	7.64986	7.76372
0.7	5.71957	5.97173	5.97137	5.92350	5.98565
0.8	3.34414	3.40518	3.40518	3.39253	3.40883
0.9	1.00000	1.00000	1.00000	1.00000	1.00000
1.0	0.00000	0.00000	0.00000	0.00000	0.00000

Table 3: The fermion determinant versus the harmonic parts ($h_1 = h_2$), on the 8×8 lattice. In each column, the normalization constant is chosen such that $\det(D) = 1$ for $h_1 = h_2 = 0.1$. Note that the fermion determinants satisfy the symmetries of the exact solution: (i) periodicities in h_1 and h_2 with period 1; (ii) reflection symmetries with respect to $h_1 = 1/2$ and $h_2 = 1/2$.

errors. The relevant question is how to minimize the dependence on these two extra parameters. More importantly, if the dependence on a_5 is too strong, then at nonzero a_5 , the effective 4D theory might differ significantly from the target theory, even in the limit $N_s \rightarrow \infty$. Therefore, it is desirable to eliminate a_5 completely and to preserve the chiral symmetry optimally for any N_s .

In the limit $N_s \rightarrow \infty$ followed by $a_5 \rightarrow 0$, the fermion determinant (including the contribution of pseudofermions) of the conventional DWF gives the effective 4D lattice Dirac operator

$$D(m_q) = (D_c + m_q)(1 + rD_c)^{-1} = m_q + (1 - rm_q)D$$

where

$$\begin{aligned}
D_c &= 2m_0 \frac{1 + \gamma_5 H_w (H_w^2)^{-1/2}}{1 - \gamma_5 H_w (H_w^2)^{-1/2}} \\
D &= D_c (1 + rD_c)^{-1} = m_0 [1 + \gamma_5 H_w (H_w^2)^{-1/2}] ,
\end{aligned}$$

while the quark propagator in background gauge field is

$$\langle q(x) \bar{q}(y) \rangle = (D_c + m_q)_{x,y}^{-1} . \quad (104)$$

Since the optimal rational approximation for $\text{sgn}(H_w) = H_w (H_w^2)^{-1/2}$ is $H_w R_Z(H_w^2)$, where $R_Z(x)$ is the Zolotarev optimal rational polynomial for

$x^{-1/2}$ (i.e., $R_Z^{(n-1,n)}(x)$ for $N_s = 2n$, or $R_Z^{(n,n)}(x)$ for $N_s = 2n + 1$), the relevant question is whether one can construct a DWF action such that its effective lattice Dirac operator for finite N_s is

$$D(m_q) = m_q + (m_0 - m_q/2)[1 + \gamma_5 H_w R_Z(H_w^2)] ,$$

in view of that the S operator in the conventional DWF is a rational polynomial approximation of $\text{sgn}(H)$. The solution to this problem is the original optimal DWF action [5]. The salient feature of the optimal DWF is that its weights $\{\omega_s\}$ (7) not only give S_{opt} , the optimal rational approximation of $\text{sgn}(H_w)$, but also eliminate a_5 completely.

However, the correlation function of $q(x)$ and $\bar{q}(y)$ is *not* equal to the quark propagator $(D_c + m_q)_{x,y}^{-1}$, nor the inverse of the effective 4D lattice Dirac operator, $D(m_q)_{x,y}^{-1}$. In other words, either $(D_c + m_q)_{x,y}^{-1}$ or $D(m_q)_{x,y}^{-1}$ *cannot* be expressed in terms of the correlation function of *local* quark fields $q(x)$ and $\bar{q}(y)$, but $q(x)$ and some “nonlocal” field variable $\bar{Q}(y)$ which depends on $\bar{q}(x)$ at all x . This posed a rather serious problem to the optimal DWF, since it implies that the basic concept of *local* fields $q(x)$ and $\bar{q}(x)$ in quantum field theory has to be abandoned in the optimal DWF. Even if $\bar{Q}(y)$ might be exponentially-local, it is still unsatisfactory in comparison with the exactly local quarks fields q and $\bar{q}$ in the conventional DWF.

The solution to this problem is presented in this paper: Appending two boundary layers at $s = 0$ and $s = N_s + 1$ with $\omega_0 = \omega_{N_s+1} = 0$. Then the correlation function of (local) quark fields $q(x)$ and $\bar{q}(y)$ constructed from these new boundary modes (37)-(38) satisfies (104). Consequently, one can derive the generating functional for n -point Green’s function of quark fields (with local Grassman sources J and $\bar{J}$)

$$e^{W[J,\bar{J}]} = \frac{\int [dU] e^{-A_g} \det[(D_c + m_q)(1 + rD_c)^{-1}] \exp \left\{ \bar{J}(D_c + m_q)^{-1} J \right\}}{\int [dU] e^{-A_g} \det[(D_c + m_q)(1 + rD_c)^{-1}]} .$$

Then the bilinear (and higher order) quark operators manifest their continuum discrete symmetries at finite lattice spacing, since the chiral projectors for both q and $\bar{q}$ are the usual $P_{\pm} = (1 \pm \gamma_5)/2$. In particular, the V-A structure of the left-handed quark current in the effective weak Hamiltonian is preserved exactly at any finite lattice spacing, which is crucial for investigating the weak matrix elements in lattice QCD, e.g., the CP violation in the $K \rightarrow \pi\pi$ decay through $\bar{K}^0$ - K^0 mixing.

From the 2D tests in Section 3, we see that the fermion determinant (ratio) of the effective lattice Dirac operator D of the conventional DWF does not agree with the exact solution for any topologically nontrivial background gauge field, even though D can capture the topology of the gauge background (i.e, reproducing topological zero and nonzero real eigenmodes with definite chirality, satisfying the Atiyah-Singer index theorem on a finite lattice, and its

axial anomaly in agreement with the topological charge density). It is interesting to see that a lattice Dirac operator satisfies the Ginsparg-Wilson relation, γ_5 -hermiticity, doubler-free, exponential-locality, and the correct continuum (free fermion) behavior, but could not reproduce the proper fermion determinant for topologically nontrivial gauge backgrounds, due to its dependence on $a_5 (\neq 0)$. This warrants further studies on a_5 effects in lattice QCD with the conventional DWF (e.g., to investigate to what extent the pion mass or other observables depending on a_5).

Finally, it is instructive to compare the quark fields in lattice QCD with Ginsparg-Wilson Dirac operator to the exactly local quark fields in the (optimal/conventional) DWF. In general, the quark action of lattice QCD with Ginsparg-Wilson Dirac operator reads

$$\mathcal{A}_q = \sum_{x,y} \bar{q}(x) D_{x,y}(m_q) q(y) , \quad (105)$$

where

$$D(m_q) = (D_c + m_q)(1 + rD_c)^{-1} = m_q + (1 - rm_q)D, \quad D = D_c(1 + rD_c)^{-1} ,$$

and D satisfies the Ginsparg-Wilson relation (29). Here we assume that D is doubler-free, exponentially-local for sufficiently smooth background gauge fields, and has the correct continuum (free fermion) behavior. Then the quark propagator in background gauge field is

$$\langle q(x) \bar{q}(y) \rangle = \frac{\int [d\bar{q}][dq] q(x) \bar{q}(y) e^{-\mathcal{A}_q}}{\int [d\bar{q}][dq] e^{-\mathcal{A}_q}} = D_{x,y}^{-1}(m_q) . \quad (106)$$

But this cannot be the valence quark propagator since it breaks the chiral symmetry even in the massless ($m_q = 0$) limit. Thus the quark propagator has to be extracted as

$$(D_c + m_q)_{x,y}^{-1} = (1 - rm_q)^{-1} [D_{x,y}^{-1}(m_q) - r\delta_{x,y}] .$$

In other words, one has to replace either $q(x)$ or $\bar{q}(y)$ in the quark bilinear of (106) with one of the followings:

$$q(x) \rightarrow \hat{q}(x) = \sum_z (1 + rD_c)_{x,z}^{-1} q(z) = \sum_z (1 - rD)_{x,z} q(z) ,$$

or

$$\bar{q}(y) \rightarrow \hat{\bar{q}}(y) = \sum_z \bar{q}(z) (1 + rD_c)_{z,y}^{-1} = \sum_z \bar{q}(z) (1 - rD)_{z,y} .$$

Note that both above transformations are exponentially-local. Suppose one replaces $q(x)$ with $\hat{q}(x)$, then the quark propagator (in background gauge field) becomes

$$\langle \hat{q}(x) \bar{q}(y) \rangle = (D_c + m_q)_{x,y}^{-1} . \quad (107)$$

Even though the concept of local field is only violated exponentially by $\hat{q}(x)$, it is still unsatisfactory from the theoretical viewpoint, in comparison with the exactly local q and $\bar{q}$ in the (optimal/conventional) DWF. To impose the exact locality of $\hat{q}$, one can rewrite the quark action as

$$\mathcal{A}_q = \sum_{x,y} \bar{q}(x)(D_c + m_q)_{x,y} \hat{q}(y) , \quad (108)$$

which, however, is also unsatisfactory since D_c is nonlocal. In general, it seems to be impossible to formulate lattice QCD with any Ginsparg-Wilson lattice Dirac operator such that the quark fields and the action both are consistent. However, if the Ginsparg-Wilson lattice Dirac operator (e.g., the overlap Dirac operator with Zolotarev rational approximation) is equal to the effective 4D lattice Dirac operator of certain DWF action (e.g., the optimal DWF), then one can resort to the DWF action, in which the quark fields q and $\bar{q}$ (e.g., (37)-(38)) are well-defined in terms of the boundary modes, with exact locality as well as the usual chiral projectors $P_{\pm} = (1 \pm \gamma_5)/2$.

Appendix

In this appendix, we derive a formula for computing the inverse of the effective 4D lattice Dirac operator, with the original optimal DWF operator (1).

From (19), we have

$$\det[\mathcal{D}(m_q)\mathcal{D}^{-1}(2m_0)] = \det[r(D_c + m_q)(1 + rD_c)^{-1}] = \det[rD(m_q)] \quad (109)$$

or equivalently,

$$\det[\mathcal{D}^{-1}(m_q)\mathcal{D}(2m_0)] = \det[r^{-1}D^{-1}(m_q)] \quad (110)$$

where $\mathcal{D}$ is the optimal DWF operator defined in (1), while $D(m_q)$ is the effective 4D lattice Dirac operator. It follows that there may exist an operator Q such that

$$\{Q\mathcal{D}^{-1}(m_q)\mathcal{D}(2m_0)Q^{-1}\}(x, 1; x', 1) = r^{-1}D^{-1}(m_q)(x, x') . \quad (111)$$

Now we want to find an explicit expression of Q . Using the Grassman integral formula

$$\frac{1}{\det\mathcal{D}(m_q)} \int [d\bar{\psi}][d\psi] \psi_s(y) \bar{\psi}_{s'}(z) e^{-\mathcal{A}_f} = \mathcal{D}^{-1}(m_q)(y, s; z, s') ,$$

we rewrite the l.h.s. of (111) as

$$\begin{aligned} & \{Q\mathcal{D}^{-1}(m_q)\mathcal{D}(2m_0)Q^{-1}\}(x, 1; x', 1) \\ &= \frac{1}{\det\mathcal{D}(m_q)} \int [d\bar{\psi}][d\psi] \{Q\psi\}(x, 1) \{\bar{\psi}\mathcal{D}(2m_0)Q^{-1}\}(x', 1) e^{-\mathcal{A}_f} . \end{aligned} \quad (112)$$

From the transformation (16)-(18), one immediately sees that the optimal DWF action (1) can be rewritten as

$$\begin{aligned}
\mathcal{A}_f &= \sum_{s,s'=1}^{N_s} \sum_{x,x'} \bar{\psi}_s(x) \mathcal{D}(m_q)(x, s; x' s') \psi_{s'}(x') \\
&= \sum_{s,s'=1}^{N_s} \sum_{x,x'} \bar{\eta}_s(x) \{(\Omega a_5 H_w - 1)^{-1} \gamma_5 \mathcal{D}(m_q) B^{-1}\}(x, s; x', s') \eta_{s'}(x') \\
&= \bar{\eta}_1(P_- - r m_q P_+) \eta_1 - \bar{\eta}_1 T_1^{-1} \eta_2 + \sum_{s=2}^{N_s-1} \left\{ \bar{\eta}_s \eta_s - \bar{\eta}_s T_s^{-1} \eta_{s+1} \right\} \\
&\quad + \bar{\eta}_{N_s} \eta_{N_s} - \bar{\eta}_{N_s} T_{N_s}^{-1} (P_+ - r m_q P_-) \eta_1
\end{aligned} \tag{113}$$

where

$$\Omega(x, s; x', s') = \omega_s \delta_{s,s'} \delta_{x,x'} ,$$

and

$$B(x, s; x', s') = \delta_{x,x'} (P_- \delta_{s,s'} + P_+ \delta_{s-1,s'}) , \tag{114}$$

$$B^{-1}(x, s; x', s') = \delta_{x,x'} (P_- \delta_{s,s'} + P_+ \delta_{s+1,s'}) , \tag{115}$$

with periodic boundary conditions in the fifth dimension.

Now it is clear that $Q = B$ would satisfy (111). This can be seen as follows. Using the transformation (16)-(18), the r.h.s. of (112) can be written as

$$\frac{\mathcal{J}}{\det \mathcal{D}(m_q)} \int [d\bar{\eta}] [d\eta] e^{-\mathcal{A}_f} \eta_1(x) \{ \bar{\eta} (\Omega a_5 H_w - 1)^{-1} \gamma_5 \mathcal{D}(2m_0) B^{-1} \}(x', 1) \tag{116}$$

where $\mathcal{J}$ is the Jacobian defined in (49). Using (113), the last expression in the integrand of (116) can be simplified as

$$\begin{aligned}
&\{ \bar{\eta} (\Omega a_5 H_w - 1)^{-1} \gamma_5 \mathcal{D}(2m_0) B^{-1} \}(x', 1) \\
&= \bar{\eta}_1(x') (P_- - P_+) - \{ \bar{\eta}_{N_s} T_{N_s}^{-1} \}(x') (P_+ - P_-) .
\end{aligned} \tag{117}$$

Thus (116) becomes

$$\frac{\mathcal{J}}{\det \mathcal{D}(m_q)} \int [d\bar{\eta}] [d\eta] e^{-\mathcal{A}_f} \left[-\eta_1(x) \bar{\eta}_1(x') - \eta_1(x) \{ \bar{\eta}_{N_s} T_{N_s}^{-1} \}(x') \right] \gamma_5 , \tag{118}$$

which can be integrated successively from $(\eta_{N_s}, \bar{\eta}_{N_s})$ to $(\eta_1, \bar{\eta}_1)$. Explicitly,

$$\begin{aligned}
&\int [d\bar{\eta}] [d\eta] e^{-\mathcal{A}_f} \eta_1(x) \{ \bar{\eta}_{N_s} T_{N_s}^{-1} \}(x') \gamma_5 \\
&= \int [d\bar{\eta}_1] [d\eta_1] e^{-\bar{\eta}_1 [(P_- - r m_q P_+) - \prod_{s=1}^{N_s} T_s^{-1} (P_+ - r m_q P_-)] \eta_1} \eta_1(x) \{ \bar{\eta}_1 \prod_{s=1}^{N_s} T_s^{-1} \}(x') \gamma_5 ,
\end{aligned}$$

which is substituted into (118) to give

$$\begin{aligned}
& \frac{\mathcal{J}}{\det \mathcal{D}(m_q)} \int [d\eta_1][d\bar{\eta}_1] e^{-\bar{\eta}_1 [(P_- - rm_q P_+) - \prod_{s=1}^{N_s} T_s^{-1} (P_+ - rm_q P_-)] \eta_1} \times \\
& \quad \eta_1(x) \left[\bar{\eta}_1 \left(-1 - \prod_{s=1}^{N_s} T_s^{-1} \right) \right] (x') \gamma_5 \\
& = \left\{ \left[(P_- - rm_q P_+) - \prod_{s=1}^{N_s} T_s^{-1} (P_+ - rm_q P_-) \right]^{-1} \left(-1 - \prod_{s=1}^{N_s} T_s^{-1} \right) \gamma_5 \right\} (x, x') \\
& = r^{-1} [(D_c + m_q)^{-1} (1 + r D_c)] (x, x') = r^{-1} D^{-1}(m_q) (x, x')
\end{aligned}$$

where (49), (19) and (52) have been used.

Therefore, we have

$$\{B \mathcal{D}^{-1}(m_q) \mathcal{D}(2m_0) B^{-1}\} (x, 1; x', 1) = r^{-1} D^{-1}(m_q) (x, x') , \quad (119)$$

and $D^{-1}(m_q)$ can be obtained by solving the linear system

$$\mathcal{D}(m_q) Y = \mathcal{D}(2m_0) B^{-1} \mathbb{1} . \quad (120)$$

The solution of (120) gives

$$D^{-1}(m_q) (x, x') = r (BY) (x, 1; x', 1) , \quad (121)$$

and the valence quark propagator (69).

Acknowledgement

This work was supported in part by National Science Council, ROC, under the grant number NSC91-2112-M002-025.

References

- [1] V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. B **125**, 136 (1983).
- [2] C. G. Callan and J. A. Harvey, Nucl. Phys. B **250**, 427 (1985).
- [3] D. B. Kaplan, Phys. Lett. B **288**, 342 (1992)
- [4] Y. Shamir, Nucl. Phys. B **406**, 90 (1993)
- [5] T. W. Chiu, Phys. Rev. Lett. **90**, 071601 (2003)
- [6] T. W. Chiu, Phys. Lett. B **552**, 97 (2003)
- [7] E. I. Zolotarev, Zap. Imp. Akad. Nauk. St. Petersburg, 30 (1877), no. 5; reprinted in his Collected works, Vol. 2, Izdat, Akad. Nauk SSSR, Moscow, 1932, p. 1-59.
- [8] N. I. Akhiezer, "Theory of approximation", Reprint of 1956 English translation, Dover, New York, 1992; "Elements of the theory of elliptic functions", Translations of Mathematical Monographs, 79, American Mathematical Society, Providence, R.I. 1990.
- [9] T. W. Chiu and S. V. Zenkin, Phys. Rev. D **59**, 074501 (1999)
- [10] P. H. Ginsparg and K. G. Wilson, Phys. Rev. D **25**, 2649 (1982)
- [11] T. W. Chiu, Phys. Rev. D **60**, 034503 (1999)
- [12] T. W. Chiu, T. H. Hsieh, C. H. Huang and T. R. Huang, Phys. Rev. D **66**, 114502 (2002)
- [13] H. Neuberger, Phys. Lett. B **417**, 141 (1998)
- [14] R. Narayanan and H. Neuberger, Nucl. Phys. B **443**, 305 (1995)
- [15] H. Neuberger, Phys. Rev. D **57**, 5417 (1998)
- [16] Y. Kikukawa and T. Noguchi, hep-lat/9902022.
- [17] R. G. Edwards and U. M. Heller, Phys. Rev. D **63**, 094505 (2001)
- [18] A. Borici, Nucl. Phys. Proc. Suppl. **83**, 771 (2000)
- [19] M. Luscher, Phys. Lett. B **428**, 342 (1998)
- [20] T. W. Chiu, Phys. Rev. D **58**, 074511 (1998)
- [21] I. Sachs and A. Wipf, Helv. Phys. Acta **65**, 652 (1992).
- [22] T. W. Chiu, hep-lat/9911010.